



Groundbreaking Insights on Penile Clamps for Managing Male Incontinence Post-Prostate Surgery

Eva Ten

Department of Urology, Franziskus Hospital, Berlin, Germany

*Corresponding Author: Eva Ten, Department of Urology, Franziskus Hospital, Berlin, Germany

Submitted: 22 Jan 2024

Accepted: 25 Jan 2024

Published: 29 Jan 2024

Citation: Eva Ten (2024). *Groundbreaking Insights on Penile Clamps for Managing Male Incontinence Post-Prostate Surgery*. *J of Clin Case Stu, Reviews & Reports*. 2(1), 1-1.

New research has surfaced with compelling findings, suggesting that penile clamps present a formidable alternative to absorbents for male patients recovering from prostate surgery, particularly those grappling with prostate cancer.

This innovative solution boasts several advantages, including effective moisture control, heightened mobility, and an overall enhancement in skincare practices.

Despite advancements in surgical techniques for prostate cancer removal, incontinence remains a prevalent issue. Residual stress incontinence, though temporary and moderate, persists as a common challenge.

Furthermore, patients may contend with urgency incontinence, a consequence of treatments such as radiotherapy or hormone therapy.

The study's revelations affirm the high efficacy of these external devices in managing stress and mixed incontinence. However, the same cannot be claimed for pure urgency incontinence, as the devices don't address the underlying urge to urinate.

Noteworthy is the almost negligible occurrence of side effects associated with these clamps, provided users adhere to the recommended conditions and maximum usage times outlined by manufacturers.

One standout device, distinguished by its flexibility and soft materials, is a closure-free clamp known as Uriclak. Notably, it emerged as the preferred choice among participants in the study.

Lead researcher Eva Ten, from the Department of Urology at Franziskus Hospital in Berlin, has spearheaded this insightful exploration into the transformative potential of these devices in post-prostate surgery care 1-[8].

References

1. Eva Ten (2023) Comparative analysis of conventional penile clamps and Uriclak device in managing male incontinence following radical, turp, or laser prostate surgery. Franziskus Hospital, Berlin. *Front. Surg.Sec. Genitourinary Surgery* Volume 10. <https://doi.org/10.3389/fsurg.2023.1301353>.
2. Lemmens JH, Broadbridge J, Macaulay M, Rees RW, Archer M, et al. (2019) Tissue response to applied loading using different designs of penile compression clamps. Published online June 27, 2019. University of Southampton, School of Health Sciences, Southampton, SO17 1BJ, UK. Department of Urology, University Hospital Southampton NHS Foundation Trust, Southampton, SO16 6YD, UK. Bristol Urological Institute, Southmead Hospital, Bristol, BS10 5NB, UK. Faculty of Nursing, University of Alberta, Edmonton, Alberta, T6G 1C9, Canada 12: 235-243.
3. Lanza EG, Contijoch MG, Miranda JEB (2009) Physiotherapy and palliative management of urinary incontinence in prostate cancer. Start point and end of the road. *Arch Esp Urol* 62: 889-895. doi: 10.4321/s0004-06142009001000014.
4. Macaulay M, Broadbridge J, Gage H, Williams P, Birch B, et al. () A trial of devices for urinary incontinence after treatment for prostate cancer. Continence and skin technology group, University College London, London, *Department of Economics, ‡ Department of Mathematics, University of Surrey, Guildford, † Continence and Skin Technology Group, University of Southampton, Southampton, § Department of Urology, University Hospital Southampton NHS Foundation Trust, Southampton, UK, and University of Alberta, Edmonton, AB, Canada 116: 432-42.
5. Anderson CA, Omar MI, Campbell SE, Hunter KF, Cody JD, et al. (2015) Conservative management of postprostatectomy urinary incontinence. *Cochrane Database Syst Rev* 1: CD001843. doi:10.1002/14651858.CD001843.pub5.
6. Fujisaki Y, Otsuka I, Kobayashi T, Miyake N, Ito K, et al. (2023) Use of the anterior prostatic urethral mucosa preservation technique during holmium laser enucleation of the prostate can reduce postoperative stress urinary incontinence. *Asian J Endosc Surg*. First published. doi: 10.1111/ases.13256.
7. Moore KN, Schieman S, Ackerman T, Dzus HY, Metcalfe JB, et al. (2004) Assessing comfort, safety, and patient satisfaction with three commonly used penile compression devices. *Urology* 63: 150-154. doi:10.1016/j.urology.2003.08.034.
8. Manufacturer's websites for male incontinence clamps. Available at: <https://www.uriclak.com/en> ; <https://patient.uwhealth.org/healthfacts/7909>.